Richard M. Marks,M.D., FACS
Professor, Orthopaedic Surgery
Director, Division of Foot/Ankle Surgery
Department of Orthopaedic Surgery

Pre-Operative Preparation

Insurance

The department of Orthopaedics will contact your insurance company to preauthorize your surgery; however, it is your responsibility to check your out-of-pocket costs such as co-pays, deductibles, or out-of network charges.

Richard M. Marks,M.D., FACS
Professor, Orthopaedic Surgery
Director, Division of Foot/Ankle Surgery
Department of Orthopaedic Surgery

Home Medications

Your current medications should be reviewed at your pre-operative physical; performed at our Pre-Admission Testing clinic or by your primary care physician.

***STOP** the following medications 7 days prior to surgery:

1. Aspirin – unless told differently by your physician or Cardiologist (patients with cardiac stents, carotid stents or coronary artery disease do **not** stop aspirin prior to surgery)
2. Anti-inflammatories (eg. Aleve, Motrin, Advil, Ibuprofen)
3. All herbal medications
4. Vitamin E
5. Multivitamins

*If you are a rheumatoid arthritis patient, it may be necessary to stop several medications 2 weeks prior to surgery. You will need to coordinate this with your rheumatologist.

*If you have any type of fusion or screws placed during your surgery, you should avoid all anti-inflammatories for six weeks after surgery.

Please call the nurse at 414-805-7442 if you develop any cold or flu symptoms prior to surgery.

NIGHT BEFORE SURGERY

DO NOT EAT OR DRINK anything after midnight. This includes, gum, mints, hard candy, tea, coffee, and water.

MORNING OF SURGERY

1. Do **NOT** use tobacco
2. Removal of all body piercings
3. Leave all valuables at home
4. Remove all nail polish
5. Do **NOT** wear make-up
6. Wear comfortable clothes with wide leg openings
7. Bring your post-op shoe if provided
8. Bring a case for your glasses or contacts